

Galaxy Formation Simulations: Problems and Answers




Neal Katz

UMass Astronomy


L. Yang, R. Dave, S. Huang, E. Scannapieco, D. Weinberg

May 2023




Implications of Baryon Cycling


- Milky Way galaxy at $z = 2.5$: 1/2 the halo mass and 1/20 the stellar mass.
- Galactic gas content is a delicate balance of inflowing gas, outflowing gas, and star formation.
- After $z \approx 2$ most inflowing gas comes from recycled winds.
- Any minor physical or numerical thing that affects this delicate balance probably will change the gas content, and hence the stellar content.




- Cold/Hot accretion
- Galactic winds
- Wind reaccretion




Galaxy Stellar Mass Function




- Changing the SPH has only a minor effect at the largest masses.
- Changing the wind launch has a major effect at the largest masses.



Galaxy Stellar Mass Function




- Changing the SPH has only a minor effect at the largest masses.
- Changing the wind launch has a major effect at the largest masses.
- The Hydro technique (SPH, AMR, AREPO, GIZMO etc.) is much less important than the feedback implementation in simulations of galaxy formation.




Fake News!


- Still need scaling laws to launch the winds.
- Wind particles are individual particles and individual particles do not properly represent hydrodynamics.
- Individual particles cannot mix metals.
- The results are highly sensitive to the exact form of the subgrid wind model and how it interacts with the numerics including the hydro code and resolution.
- Why not just solve the problem by brute force?



Why not just solve the problem by brute force?


- Winds are dominated by cold gas that is thought to be entrained.
- Simulations show that the convergence of this process does not occur until the resolution is Solar System in scale and Jupiter in mass.
- Even then clouds are not accelerated.
- Entrainment does not occur unless perhaps there is magnetic draping.






More Fake News; SAD!

- Any claims that any simulation can have winds develop **naturally** are highly dubious.
- One expects that the problems would be even worse for propagating winds through the CGM into the IGM.
- Cold gas clouds traveling through a less dense, hot CGM should typically not slow down but slowly disintegrate on a time scale of many t_{cc} , which does not happen in current simulations.
- The interactions at wind/halo gas interfaces in the CGM occur on scales that are much below the resolution of any galaxy formation simulation, including FIRE, Illustris, FOGGIE, and Eagle.




More Fake News; SAD!

- Any claims that any simulation can have winds develop **naturally** are highly dubious.
- One expects that the problems would be even worse for propagating winds through the CGM into the IGM.
- Cold gas clouds traveling through a less dense, hot CGM should typically not slow down but slowly disintegrate on a time scale of many t_{cc} , which does not happen in current simulations.
- The interactions at wind/halo gas interfaces in the CGM occur on scales that are much below the resolution of any galaxy formation simulation, including FIRE, Illustris, FOGGIE, and Eagle.
- **Do not have and will not have for many years the ability to simulate superwinds leaving galaxies and in particular their interactions with the CGM and IGM so we must develop a subgrid model.**
- PhEW, there is another way forward.




Tell me what you really really want


- Want the model to be limited by our physical assumptions and not by numerics.
- Want a method that must be as independent of resolution as possible.
- Want a method that must be as independent of hydro technique as possible.
 - ◆ Works with SPH, AMR, and moving mesh codes (e.g. AREPO and GIZMO).
- Want to try to limit the number of free parameters.
- Want it to globally conserve mass, momentum, and energy.
- Most importantly: want it to correctly represent the physics and not depend on unknown numerics.



Physically Evolved Winds (PhEW)

- Wind particles are launched as before.
- They are evolved analytically using microphysics that depends on the surrounding medium.
- Wind particles are "removed" and added to their surroundings when their mass becomes small relative to the surrounding particles.






Physically Evolved Winds (PhEW)



- Cloud motion affected by:
 - ◆ gravity,
 - ◆ ram pressure.

- Cloud temperature affected by:
 - ◆ radiative and adiabatic heating and cooling,
 - ◆ ram pressure heating,
 - ◆ conduction.


- Clouds lose mass, thermal energy, and metals to surroundings owing to:
 - ◆ Kelvin-Helmholtz and Rayleigh Taylor instabilities,
 - ◆ Conductive evaporation.




Physically Evolved Winds (PhEW)




- Assume each wind particle is made of many cylindrical clouds with mass M_c , temperature T_c , uniform density, ρ_c , and radius R_c .
- Clouds create a conductive bow shock, creating a post-shock medium whose properties depend on the ambient conditions and the cloud speed.
- Assume clouds are in pressure equilibrium and thermal contact with the post-shock medium.




Physically Evolved Winds (PhEW)




- The model can be set using high resolution single cloud simulations.
- These simulations can also determine cloud absorption line properties.
- Still need parameters:
 - ◆ M_{cloud} : the sub-cloud mass,
 - ◆ f_s : the fraction of the Spitzer rate for conductive processes.
 - ◆ f_{KH} : controls the Kelvin-Helmholtz destruction time.




Galaxy Stellar Mass Function




- Proof of concept simulation; parameters have not been tuned.
- Better matches the GSMF at the knee.
- Too many high mass galaxies.




Resolution and M_c Effects on GSMFs




- The old wind model is very sensitive to resolution and hydro method.
- The PhEW model is not very sensitive to resolution or M_c or hydro method.




Anywhere the Wind Blows




- When the KHI dominates the mass loss, $\tau_{\text{KH}} \propto f_{\text{KH}} M_c^{1/3}$.
- When conductive evaporation dominates the mass loss, $\tau_{\text{ev}} \propto M_c^{2/3} / f_s$.




How Galaxies get their gas in PhEW




- Redefine wind reaccretion as fraction of particle formally in wind.
- Wind reaccretion now dominates accretion below about $10^{12.3} M_{\odot}$.
- The total amount of cold accretion is similar but the amount of hot and wind accretion increases.



CGM gas metallicities in PhEW




- Metallicity without PhEW is trimodal (40% of particles have $Z \approx 0$).
- With PhEW metallicities have a single peak around 10^{-1} .
- In PhEW the metal distributions are robust to numerical resolution.




Conclusions

- The Hydro technique is much less important than the feedback implementation in simulations of galaxy formation.
- Do not and will not have for many years the resolution to correctly model superwinds leaving galaxies and their interactions with the CGM and IGM so we must develop a subgrid model.



Conclusions

- The Hydro technique is much less important than the feedback implementation in simulations of galaxy formation.
- Do not and will not have for many years the resolution to correctly model superwinds leaving galaxies and their interactions with the CGM and IGM so we must develop a subgrid model.
- This is even more true for AGN feedback!
- Physically Evolved Winds (PhEW) are a way forward.
 - ◆ Can be tuned to match very high resolution ISM simulations.
 - ◆ Can be used in any code and is almost independent of resolution.
- Current "proof of concept" simulations look promising.



Conclusions

- The Hydro technique is much less important than the feedback implementation in simulations of galaxy formation.
- Do not and will not have for many years the resolution to correctly model superwinds leaving galaxies and their interactions with the CGM and IGM so we must develop a subgrid model.
- This is even more true for AGN feedback!
- Physically Evolved Winds (PhEW) are a way forward.
 - ◆ Can be tuned to match very high resolution ISM simulations.
 - ◆ Can be used in any code and is almost independent of resolution.
- Current "proof of concept" simulations look promising.
- Making Galaxy Formation Simulations Great Again!
- Huang, Katz et al MNRAS (2020 497 2586; 2022 509 6091).